

Persons Living with a Diagnosis of HIV
and/or AIDS Reported in HIV
Prevention Planning Lake

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by selected characteristics, HIV Prevention Planning Lake

Characteristic	Living with a diagnosis of HIV infection in 2014			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Sex							
Males	107.2	120	80%	52	78%	68	82%
Females	25.6	30	20%	15	22%	15	18%
Age at end of year							
<13	*	-	-	-	-	-	-
13-14	*	-	-	-	-	-	-
15-19	*	-	-	-	-	-	-
20-24	*	1	1%	1	1%	-	-
25-29	78.2	10	7%	8	12%	2	2%
30-34	*	3	2%	3	4%	-	-
35-39	133.7	17	11%	8	12%	9	11%
40-44	97.4	14	9%	4	6%	10	12%
45-49	182.1	29	19%	12	18%	17	20%
50-54	214.4	39	26%	15	22%	24	29%
55-64	80.9	28	19%	13	19%	15	18%
65+	21.8	9	6%	3	4%	6	7%
Race/Ethnicity							
White, not Hispanic	48.8	100	67%	43	64%	57	69%
Black/African American, not Hispanic	391.7	33	22%	15	22%	18	22%
Hispanic/Latino	109.6	10	7%	3	4%	7	8%
Asian/Pacific Islander	*	1	1%	1	1%	-	-
American Indian/Alaska Native	*	1	1%	1	1%	-	-
Unknown	*	5	3%	4	6%	1	1%
Race/Sex							
White, not Hispanic Males	85.3	85	57%	37	55%	48	58%
White, not Hispanic Females	14.2	15	10%	6	9%	9	11%
Black/African American, not Hispanic Males	539.5	22	15%	9	13%	13	16%
Black/African American, not Hispanic Females	253.1	11	7%	6	9%	5	6%
Hispanic/Latino Males	162.0	8	5%	2	3%	6	7%
Hispanic/Latina Females	*	2	1%	1	1%	1	1%
Asian/Pacific Islander Males	*	1	1%	1	1%	-	-
Asian/Pacific Islander Females	*	-	-	-	-	-	-
American Indian/Alaska Native Males	*	-	-	-	-	-	-
American Indian/Alaska Native Females	*	1	1%	1	1%	-	-
Unknown	*	5	3%	4	6%	1	1%
Total	65.4	150		67		83	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Asterisk (*) indicates rate not calculated for case count < 5 due to unstable rates.

^a The rate is the number of persons living with a diagnosis of HIV infection per 100,000 population calculated using 2014 U.S. Census estimates.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by transmission category, HIV Prevention Planning Lake

Transmission Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	76	63%	31	60%	45	66%
Injection drug use (IDU)	11	9%	2	4%	9	13%
Male-to-male sexual contact and IDU	5	4%	2	4%	3	4%
Heterosexual contact	9	8%	5	10%	4	6%
Other/unknown	19	16%	12	23%	7	10%
Subtotal	120	100%	52	100%	68	100%
Female adult or adolescent						
Injection drug use	1	3%	1	7%	-	-
Heterosexual contact	19	66%	8	57%	11	73%
Other/unknown	9	31%	5	36%	4	27%
Subtotal	29	100%	14	100%	15	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	-	-
Subtotal	1	100%	1	100%	-	-
Total	150		67		83	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by current disease status and by exposure category, HIV Prevention Planning Lake

Exposure Category ^a	Living with a diagnosis of HIV infection in 2014		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male-to-male sex (MSM) only	73	49%	30	45%	43	52%
Injection drug use (IDU) only	7	5%	-	-	7	8%
Heterosexual contact only	28	19%	13	19%	15	18%
MSM & IDU	5	3%	2	3%	3	4%
IDU & Heterosexual contact	5	3%	3	4%	2	2%
MSM & Heterosexual contact	3	2%	1	1%	2	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-
Perinatal exposure	1	1%	1	1%	-	-
Other/unknown	28	19%	17	25%	11	13%
Total	150		67		83	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by transmission category, HIV Prevention Planning Lake

Transmission Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	61	72%	12	55%	2	25%	-	-	-	-	<10	*
Injection drug use (IDU)	3	4%	4	18%	4	50%	-	-	-	-	-	-
Male-to-male sexual contact and IDU	4	5%	1	5%	-	-	-	-	-	-	-	-
Heterosexual contact	5	6%	1	5%	1	13%	-	-	-	-	<10	*
Other/unknown	12	14%	4	18%	1	13%	<10	*	-	-	<10	*
Subtotal	85	100%	22	100%	8	100%	<10	*	-	-	<10	*
Female adult or adolescent												
Injection drug use	-	-	-	-	1	50%	-	-	-	-	-	-
Heterosexual contact	11	79%	7	64%	1	50%	-	-	-	-	-	-
Other/unknown	3	21%	4	36%	-	-	-	-	<10	*	<10	*
Subtotal	14	100%	11	100%	2	100%	-	-	<10	*	<10	*
Child (<13 yrs at diagnosis)												
Perinatal	-	-	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	100%	-	-	-	-	-	-	-	-	-	-
Subtotal	1	100%	-	-	-	-	-	-	-	-	-	-
Total	100		33		10		<10		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by race/ethnicity and by exposure category, HIV Prevention Planning Lake

Exposure Category ^a	Living with a Diagnosis of HIV Infection in 2014											
	White, not Hispanic		Black/African American		Hispanic/Latino		Asian/Pacific Islander		American Indian/Alaska Native		Unknown	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sex (MSM) only	60	60%	12	36%	-	-	-	-	-	-	<10	*
Injection drug use (IDU) only	2	2%	2	6%	3	30%	-	-	-	-	-	-
Heterosexual contact only	16	16%	8	24%	2	20%	-	-	-	-	<10	*
MSM & IDU	4	4%	1	3%	-	-	-	-	-	-	-	-
IDU & Heterosexual contact	1	1%	2	6%	2	20%	-	-	-	-	-	-
MSM & Heterosexual contact	1	1%	-	-	2	20%	-	-	-	-	-	-
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	1%	-	-	-	-	-	-	-	-	-	-
Other/unknown	15	15%	8	24%	1	10%	<10	*	<10	*	<10	*
Total	100		33		10		<10		<10		<10	

Notes: Living with a diagnosis of HIV infection represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Dash (-) indicates no cases were reported for the given category. Asterisk (*) indicates percent not calculated for column < 10 cases due to small numbers.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by selected characteristics, HIV Prevention Planning Lake

Living with a Diagnosis of HIV Infection

Characteristic	2010		2011		2012		2013		2014	
	No.	%								
Sex										
Males	99	78%	103	78%	107	79%	116	80%	120	80%
Females	28	22%	29	22%	29	21%	29	20%	30	20%
Age at end of year										
<13	-	-	-	-	-	-	-	-	-	-
13-14	-	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-	-
20-24	4	3%	3	2%	4	3%	1	1%	1	1%
25-29	5	4%	6	5%	5	4%	10	7%	10	7%
30-34	10	8%	8	6%	9	7%	8	6%	3	2%
35-39	15	12%	19	14%	18	13%	14	10%	17	11%
40-44	23	18%	18	14%	14	10%	17	12%	14	9%
45-49	26	20%	25	19%	27	20%	27	19%	29	19%
50-54	23	18%	28	21%	30	22%	37	26%	39	26%
55-64	19	15%	23	17%	26	19%	23	16%	28	19%
65+	2	2%	2	2%	3	2%	8	6%	9	6%
Race/Ethnicity										
White, not Hispanic	82	65%	86	65%	89	65%	95	66%	100	67%
Black/African American, not Hispanic	29	23%	30	23%	31	23%	33	23%	33	22%
Hispanic/Latino	10	8%	10	8%	10	7%	10	7%	10	7%
Asian/Pacific Islander	1	1%	1	1%	1	1%	1	1%	1	1%
American Indian/Alaska Native	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	4	3%	4	3%	4	3%	5	3%	5	3%
Race/Sex										
White, not Hispanic Males	69	54%	72	55%	75	55%	81	56%	85	57%
White, not Hispanic Females	13	10%	14	11%	14	10%	14	10%	15	10%
Black/African American, not Hispanic Males	18	14%	19	14%	20	15%	22	15%	22	15%
Black/African American, not Hispanic Females	11	9%	11	8%	11	8%	11	8%	11	7%
Hispanic/Latino Males	8	6%	8	6%	8	6%	8	6%	8	5%
Hispanic/Latina Females	2	2%	2	2%	2	1%	2	1%	2	1%
Asian/Pacific Islander Males	1	1%	1	1%	1	1%	1	1%	1	1%
Asian/Pacific Islander Females	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Males	-	-	-	-	-	-	-	-	-	-
American Indian/Alaska Native Females	1	1%	1	1%	1	1%	1	1%	1	1%
Unknown	4	3%	4	3%	4	3%	5	3%	5	3%
Total	127		132		136		145		150	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by transmission category, HIV Prevention Planning Lake

Living with a Diagnosis of HIV Infection

Transmission Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male adult or adolescent										
Male-to-male sexual contact	59	60%	63	61%	67	63%	73	63%	76	63%
Injection drug use (IDU)	11	11%	11	11%	11	10%	11	9%	11	9%
Male-to-male sexual contact and IDU	5	5%	5	5%	5	5%	5	4%	5	4%
Heterosexual contact	9	9%	9	9%	9	8%	9	8%	9	8%
Other/unknown	15	15%	15	15%	15	14%	18	16%	19	16%
Subtotal	99	100%	103	100%	107	100%	116	100%	120	100%
Female adult or adolescent										
Injection drug use	1	4%	1	4%	1	4%	1	4%	1	3%
Heterosexual contact	19	70%	19	68%	19	68%	19	68%	19	66%
Other/unknown	7	26%	8	29%	8	29%	8	29%	9	31%
Subtotal	27	100%	28	100%	28	100%	28	100%	29	100%
Child (<13 yrs at diagnosis)										
Perinatal	-	-	-	-	-	-	-	-	-	-
Other/unknown	1	100%	1	100%	1	100%	1	100%	1	100%
Subtotal	1	100%	1	100%	1	100%	1	100%	1	100%
Total	127		132		136		145		150	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aTransmission categories are mutually exclusive, hierarchical risk categories determined by the CDC. A person with multiple risks is only represented in the highest category based on the CDC hierarchical algorithm.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.

Reported persons living with a diagnosis of HIV infection as of June 30, 2015, by year and by exposure category, HIV Prevention Planning Lake

Living with a Diagnosis of HIV Infection

Exposure Category ^a	2010		2011		2012		2013		2014	
	No.	%								
Male-to-male sex (MSM) only	56	44%	60	45%	64	47%	70	48%	73	49%
Injection drug use (IDU) only	7	6%	7	5%	7	5%	7	5%	7	5%
Heterosexual contact only	28	22%	28	21%	28	21%	28	19%	28	19%
MSM & IDU	5	4%	5	4%	5	4%	5	3%	5	3%
IDU & Heterosexual contact	5	4%	5	4%	5	4%	5	3%	5	3%
MSM & Heterosexual contact	3	2%	3	2%	3	2%	3	2%	3	2%
MSM & IDU & Heterosexual contact	-	-	-	-	-	-	-	-	-	-
Perinatal exposure	1	1%	1	1%	1	1%	1	1%	1	1%
Other/unknown	22	17%	23	17%	23	17%	26	18%	28	19%
Total	127		132		136		145		150	

Notes: Living with a diagnosis of HIV infection by year (2010-2014) represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of December 31, 2014. Persons living with a diagnosis of HIV infection represent persons living in Ohio as of December 31, 2014, regardless of whether or not the person was a resident of Ohio at time of initial HIV and/or AIDS diagnosis.

^aExposure categories are mutually exclusive risk categories. All possible combinations of risks are represented among exposure categories. A person with multiple risks is represented in the exposure category identifying all the reported ways in which that person may have been exposed to HIV.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through December 31, 2014.